

Corticotropin-releasing factor increases mouse ventral tegmental area dopamine neuron firing through a protein kinase C-dependent enhancement of I_h

M. J. Wanat^{1,2}, F. W. Hopf², G. D. Stuber², P. E. M. Phillips³ and A. Bonci^{2,4}

¹Neuroscience Graduate Program at University of California, San Francisco, San Francisco, CA, USA

²Ernest Gallo Clinic and Research Center, Emeryville, CA, USA

³Department of Psychiatry & Behavioural Sciences and Pharmacology, University of Washington, Seattle, WA, USA

⁴Department of Neurology and Wheeler Center for the Neurobiology of Addiction, University of California, San Francisco, San Francisco, CA, USA

Stress induces the release of the peptide corticotropin-releasing factor (CRF) into the ventral tegmental area (VTA), and also increases dopamine levels in brain regions receiving dense VTA input. Therefore, stress may activate the mesolimbic dopamine system in part through the actions of CRF in the VTA. Here, we explored the mechanism by which CRF affects VTA dopamine neuron firing. Using patch-clamp recordings from brain slices we first determined that the presence of I_h is an excellent predictor of dopamine content in mice. We next showed that CRF dose-dependently increased VTA dopamine neuron firing, which was prevented by antagonism of the CRF receptor-1 (CRF-R1), and was mimicked by CRF-R1 agonists. Inhibition of the phospholipase C (PLC)–protein kinase C (PKC) signalling pathway, but not the cAMP–protein kinase A (PKA) signalling pathway, prevented the increase in dopamine neuron firing by CRF. Furthermore, the effect of CRF on VTA dopamine neurons was not attenuated by blockade of I_A , $I_{K(Ca)}$ or I_{Kir} , but was completely eliminated by inhibition of I_h . Although cAMP-dependent modulation of I_h through changes in the voltage dependence of activation is well established, we surprisingly found that CRF, through a PKC-dependent mechanism, enhanced I_h independent of changes in the voltage dependence of activation. Thus, our results demonstrated that CRF acted on the CRF-R1 to stimulate the PLC–PKC signalling pathway, which in turn enhanced I_h to increase VTA dopamine neuron firing. These findings provide a cellular mechanism of the interaction between CRF and dopamine, which can be involved in promoting the avoidance of threatening stimuli, the pursuit of appetitive behaviours, as well as various psychiatric conditions.

(Received 17 December 2007; accepted after revision 25 February 2008; first published online 28 February 2008)

Corresponding author A. Bonci: Ernest Gallo Clinic and Research Center, 5858 Horton St, Suite 200, Emeryville, CA 94608, USA. Email: antonello.bonci@ucsf.edu

Stressful stimuli induce the release of corticotropin-releasing factor (CRF) from the hypothalamus, which in turn initiates the canonical stress response pathway (Sarnyai *et al.* 2001). CRF exerts its cellular effects by activating one of its two known G-protein-coupled receptors, the CRF-receptor 1 and 2 (CRF-R1 and CRF-R2) (Hauger *et al.* 2006). CRF receptor stimulation primarily activates the cAMP–protein kinase A (PKA) signalling cascade in neuronal systems (Haug & Storm, 2000; Jedema & Grace, 2004; Hauger *et al.* 2006), though CRF receptors also can couple to the phospholipase C (PLC)–protein kinase C (PKC) pathway (Ungless *et al.* 2003; Tan *et al.* 2004; Hauger *et al.* 2006). Stress not only

increases glucocorticoid levels through the cellular actions of CRF in the hypothalamic–pituitary–adrenal axis, but also stimulates the release of CRF in extra-hypothalamic brain regions, such as the ventral tegmental area (VTA) (Wang *et al.* 2005; Hauger *et al.* 2006).

The VTA and dopamine system is involved with numerous behaviours including motor activity (Beninger, 1983; Zhou & Palmiter, 1995), motivation (Salamone, 1996; Wise, 2004; Phillips *et al.* 2007) and drug-seeking (Kauer, 2004; McFarland *et al.* 2004). Interestingly, the dopamine system is also directly activated by stressful stimuli (Horger & Roth, 1996), as restraint stress increases the firing rate of putative dopamine neurons (Anstrom & Woodward, 2005). In addition, foot-shock, handling, tail-flick and social defeat stressors all increase dopamine levels in brain regions receiving dense input from the

This paper has online supplemental material.

VTA including the prefrontal cortex, nucleus accumbens and amygdala (Abercrombie *et al.* 1989; Tidey & Miczek, 1996; Inglis & Moghaddam, 1999; McFarland *et al.* 2004). Both stress and increased dopamine levels can improve the execution of context-dependent behaviours (Rose, 1987; Blanchard *et al.* 1999; Cagniard *et al.* 2006), which suggests that moderate stress-induced activation of the dopamine system can enhance performance on specific tasks. In addition, CRF receptor activation enhances the performance of appetitive- and dopamine-dependent tasks (Pecina *et al.* 2006; Yang *et al.* 2006), suggesting that stress may affect the dopamine system in part through the actions of CRF.

Further highlighting the interaction between CRF and the dopamine system, intraventricular injections of CRF dose-dependently increase dopamine levels in the prefrontal cortex (Lavicky & Dunn, 1993), while CRF receptor antagonists reduce cocaine-induced dopamine overflow in the nucleus accumbens (Lodge & Grace, 2005). Additionally, intra-VTA application of CRF stimulates dopamine-related behaviours such as drug-seeking (Wang *et al.* 2005) and locomotor activity (Kalivas *et al.* 1987). Given the abundant evidence suggesting an interaction between the dopamine system and CRF, there is a surprising paucity of information regarding the cellular effect of CRF on VTA dopamine neurons. In other brain regions, CRF can increase neuronal firing rate through activation of the cAMP-PKA pathway (Aldenhoff *et al.* 1983; Haug & Storm, 2000; Jedema & Grace, 2004), and can act on a number of ionic conductances including $I_{K(Ca)}$ (Aldenhoff *et al.* 1983), I_{Kir} (Kuryshv *et al.* 1997) and I_h (Qiu *et al.* 2005). Also, CRF potentiates NMDA receptor currents on VTA dopamine neurons (Ungless *et al.* 2003), but how CRF affects the firing of these neurons is unknown. Since stress can stimulate the release of CRF in the VTA (Wang *et al.* 2005), increase putative VTA dopamine neuron firing (Anstrom & Woodward, 2005) and induce dopamine release in brain regions receiving VTA input (Abercrombie *et al.* 1989; Tidey & Miczek, 1996; Inglis & Moghaddam, 1999; McFarland *et al.* 2004), we hypothesized that CRF would directly excite VTA dopamine neuron firing. In the current study, we show that CRF acted on the CRF-R1 to stimulate the PLC-PKC signalling pathway, which enhanced I_h and increased the firing rate in VTA dopamine neurons.

Methods

Electrophysiology

All procedures conformed to animal care standards set forth by the National Institute of Health and the Ernest Gallo Clinic and Research Center (EGCRC). Horizontal VTA brain slices from 3- to 5-week-old C57BL/6 mice (Charles River, Hollister, CA, USA) were prepared as

previously described (Ungless *et al.* 2003). CRF receptor transgenic mice were generously provided by Wylie Vale (Scripps Research Institute, La Jolla, CA, USA) and were bred at the EGCRC. Rodents were anaesthetized with halothane and immediately decapitated. All solutions used were saturated with 95% O₂-5% CO₂. Brain slices 170 μ m thick were cut in a chilled solution that contained (mM): 87 NaCl, 2.5 KCl, 1.25 NaH₂PO₄, 25 NaHCO₃, 0.5 CaCl₂, 7 MgCl₂ and 75 sucrose. Slices recovered for ~1 h at 32°C in an artificial cerebral spinal fluid (aCSF), with 295–305 osmolarity, and contained (mM): 126 NaCl, 2.5 KCl, 1.1 NaH₂PO₄, 1.4 MgCl₂, 2.4 CaCl₂, 11 D-glucose and 26 NaHCO₃. Picrotoxin (100 μ M) was added to the aCSF before recordings to block GABA_A input on recorded neurons.

Whole-cell patch-clamp recordings with 2–6 M Ω electrodes were made with an Axopatch 1D amplifier using Clampex 8.0 (Axon Instruments, Union City, CA, USA) and Igor Pro (Wavemetrics, Lake Oswego, OR, USA) as data acquisition programs. VTA neurons included in this study were located immediately medial to the medial terminal nucleus of the accessory optic tract. Stable firing neurons (~10 min baseline) for long-lasting recordings were found only deep in the tissue, and were recorded in current-clamp mode using an internal recording solution of 130 mM KOH, 105 mM methanesulphonic acid, 17 mM HCl, 20 mM Hepes, 0.2 mM EGTA, 2.8 mM NaCl, 2.5 mg ml⁻¹ Mg-ATP and 0.25 mg ml⁻¹ Mg-GTP. We attempted voltage-clamp recordings using this internal solution, but were unable to get stable recordings over time; thus, we used a caesium-based internal solution containing: 117 mM caesium methanesulphonate, 20 mM Hepes, 0.4 mM EGTA, 2.8 mM NaCl, 5 mM TEA-Cl, 2.5 mg ml⁻¹ Mg-ATP and 0.25 mg ml⁻¹ Mg-GTP. Internal recording solutions used were at pH = 7.2–7.4 and at an osmolarity between 275 and 285. For immunohistochemical-staining experiments, 1.0% biocytin was included in the recording solution. Neurons were visualized with an upright microscope using infrared differential interference contrast illumination. aCSF at 32–34°C was continuously perfused at 2.0–2.5 ml min⁻¹ over brain slices. Brain slices from a minimum of three mice were used for each treatment. Neurons from control and drug treatment groups were from different populations (e.g. not from the same neuron). All drugs were bath applied unless otherwise specified. Only one experiment would be done on a single neuron from a given brain slice. Finally, qualitatively similar CRF-induced increases in dopamine neuron firing were found using cell-attached recordings (see online Supplemental Fig. 1), in agreement with the finding from others in rats (Korotkova *et al.* 2006). However, all experiments in the study utilized whole-cell recordings, since this recording technique allowed us to confidently identify dopamine neurons by the detection of I_h under

voltage-clamp (Fig. 1A). In addition, with whole-cell recording techniques, one can directly apply intracellular pharmacological agents to the recorded neuron for the determination of cell-autonomous effects.

Immunocytochemistry

For immunohistochemical staining, brain slices were fixed in 4% formaldehyde for 2 h. Slices were washed with phosphate-buffered saline (PBS) and refrigerated until pre-blocking the tissue with PBT (0.2% triton, bovine serum albumin, 0.2 g (100 ml)⁻¹ in PBS) and 5% normal goat serum at 25°C for 2 h on a shaker. Slices incubated at 4°C for 2 days with 1 : 100 rabbit anti-tyrosine hydroxylase (Chemicon, Temecula, CA, USA) and then washed with PBT. Finally, slices incubated overnight at 4°C with 1 : 50 FITC-anti-rabbit and 6.5 μ l ml⁻¹ Texas red conjugated streptavidin (Jackson ImmunoResearch, West Grove, PA, USA) in PBT, then washed, mounted and visualized with a Zeiss LSM 510 META confocal microscope.

Reagents and statistical analysis

All drugs were obtained from Sigma (St Louis, MO, USA), except human/rat CRF (Sigma and Bachem, Torrance, CA, USA), anti-sauvagine-30 (Polypeptide Laboratories), ZD-7288 (Tocris, Ellisville, MO, USA), PDBU (Calbiochem, San Diego, CA, USA), CP-154,56 (generous gift from Pfizer), and ovine CRF, d-Phe-CRF, urocortin II and CRF 6-33 (all from Bachem, Torrance, CA, USA). ZD-7288 was dissolved in aCSF. All other drugs were dissolved in DMSO at a final concentration of less than 0.1% and then added to aCSF for experiments. The firing rate was determined in 10 s sweeps and averaged into 5 min bins for statistical analysis. All values are expressed as mean \pm s.e.m. Unless otherwise noted, statistical significance was assessed using a two-tailed unpaired Student's *t* test.

Results

To examine how CRF affects VTA dopamine neuron firing, we made whole-cell patch-clamp recordings from spontaneously firing neurons from mouse brain slices (baseline firing 1.90 ± 0.05 Hz, $n = 192$). The presence of the hyperpolarization-activated, cyclic nucleotide-regulated cation current (I_h) is used as an indicator of dopamine content (Grace & Onn, 1989; Cameron *et al.* 1997; Ford *et al.* 2006); however, a recent study has questioned the validity of this link in the rat VTA (Margolis *et al.* 2006). We examined whether I_h predicted dopamine content in mice by using immunocytochemistry to assay for the presence of tyrosine hydroxylase (TH), the rate-limiting enzyme in dopamine synthesis. In contrast

to the recent findings in the rat (Margolis *et al.* 2006), 98% of neurons (53/54) exhibiting an I_h also contained TH, indicating that I_h is an excellent predictor of dopamine content in mice (Fig. 1A). In a subset of these neurons, we examined whether application of a dopamine D2 receptor agonist would hyperpolarize the VTA neuron, which is indicative of an auto-inhibitory response previously reported in dopamine neurons (Lacey *et al.* 1987; Johnson & North, 1992). Indeed, we found that quinpirole (3 μ M) hyperpolarized 7/7 VTA neurons (-7.2 ± 1.5 mV) that exhibited I_h and contained TH immunoreactivity, which suggests that hyperpolarization by D2 receptor agonists can also predict dopamine content in mouse VTA slices. For all subsequent experiments, we recorded only from neurons with I_h , since they presumably were dopaminergic.

Application of CRF (1 μ M, 10 min), which potentiates glutamate receptor currents in the VTA (Ungless *et al.* 2003), also increased the firing rate of dopamine neurons (Fig. 1B). Although the time course of the CRF effect was variable with some neurons exhibiting a decrement in the CRF-induced increase in firing during CRF application (Fig. 1B), while other neurons exhibited a prolonged increase in firing even after removal of CRF (see Supplemental Fig. 2), the maximal increase in firing rate was consistent across neurons. On average, CRF elicited a $37.6 \pm 5.1\%$ increase over baseline firing ($n = 14$, $P < 0.001$, Fig. 1B and C). The CRF-induced increase in firing was concentration dependent, as 500 nM CRF significantly increased the firing by $21.5 \pm 7.5\%$ over baseline ($n = 6$, $P < 0.01$, Fig. 1D), while 100 nM CRF application was without effect on the firing rate ($3.8 \pm 2.6\%$ over baseline, $n = 5$, $P > 0.05$, Fig. 1D). All further experiments used CRF at a concentration of 1 μ M.

Either the CRF-R1 or the CRF-R2 receptor subtype could mediate the excitatory effect of CRF on VTA dopamine neurons, as both are present in the VTA (Van Pett *et al.* 2000; Ungless *et al.* 2003). The CRF-R1 agonist, ovine CRF (oCRF, 1 μ M), significantly increased the firing rate ($29.6 \pm 5.6\%$ over baseline, $n = 7$, $P < 0.001$, Fig. 2A and D), while the CRF-R2 agonist, Urocortin II (UCN II, 1 μ M), was without effect ($3.4 \pm 3.0\%$ over baseline, $n = 7$, $P > 0.05$, Fig. 2B and D), suggesting a primary role for the CRF-R1 in the CRF enhancement of VTA dopamine neuron firing. In agreement, 3 μ M CP-154,526, a CRF-R1 antagonist, significantly reduced the maximal effect of CRF on the firing rate ($15.9 \pm 4.6\%$ over baseline, $n = 7$, $P < 0.05$ compared to CRF alone, Fig. 2C and D) as did 1 μ M d-Phe-CRF, a non-specific CRF receptor antagonist ($9.5 \pm 5.8\%$ over baseline, $n = 7$, $P < 0.01$ compared to CRF alone, Fig. 2C and D). In contrast, 250 nM anti-sauvagine-30 (AS-30), a CRF-R2 antagonist, did not prevent the CRF-induced increase in VTA dopamine neuron firing ($30.4 \pm 7.3\%$ over baseline, $n = 7$, $P > 0.05$ compared to CRF alone, Fig. 2C and D).

We next examined the effect of CRF on dopamine neuron firing in mice deficient for CRF-R1 or CRF-R2 to unequivocally demonstrate a role for CRF-R1 in mediating this effect. In CRF-R1^{+/+} mice, CRF robustly increased the firing (38.6 ± 6.1% over baseline, $n = 5$, $P < 0.001$, Fig. 2E), while in the CRF-R1^{+/-} mice, CRF induced a more modest, though significant enhancement in the firing rate (19.1 ± 6.6% over baseline, $n = 6$, $P < 0.01$, Fig. 2E). Importantly, CRF had no effect on the firing rate in the CRF-R1^{-/-} mice (5.9 ± 7.2% over baseline, $n = 4$, $P < 0.05$ relative to CRF-R1^{+/+}, Fig. 2E), which indicates a critical role of the CRF-R1. Also in agreement with our pharmacological experiments suggesting no role for CRF-R2, CRF augmented the baseline firing to similar levels in the CRF-R2^{-/-} mice (26.0 ± 5.5% over baseline, $n = 7$, Fig. 2F), the CRF-R2^{+/-} mice (19.6 ± 6.5% over baseline, $n = 10$, Fig. 2F) and the CRF-R2^{+/+} mice (22.9 ± 3.2% over baseline, $n = 5$, Fig. 2F). Finally, the

CRF-binding protein (CRF-BP) can also participate in the cellular actions of CRF in the VTA (Ungless *et al.* 2003). However, the CRF-BP antagonist CRF 6-33 (1 μM) did not alter the effect of CRF on the firing rate (33.0 ± 5.7% over baseline, $n = 6$, $P > 0.05$ compared to CRF alone, data not shown). Thus, both pharmacological and transgenic methods support a role for CRF-R1, but not CRF-R2 or CRF-BP, in the CRF-mediated enhancement of dopamine neuron firing.

In order to identify the intracellular signalling pathway activated by CRF which increases VTA dopamine neuron firing, we included pathway-specific inhibitors in the intracellular recording solution. Although CRF receptors predominately couple to the cAMP-PKA pathway (Hauger *et al.* 2006), CRF surprisingly still increased dopamine neuron firing when this pathway was blocked with 100 μM Rp-cAMPs, a cAMP analogue that inhibits PKA activity (34.4 ± 11.1% over baseline, $n = 8$, $P > 0.05$ relative to




Figure 1. CRF increased the firing rate of VTA dopamine neurons in the mouse

The presence of I_h predicted dopamine content in mouse VTA neurons. A1, example neuron where a 250 ms hyperpolarizing voltage step from -60 mV to -120 mV elicited a slowly developing inward current. The magnitude of I_h was calculated by subtracting the instantaneous current (IS) from the steady-state current (SS) achieved during the voltage step. Scale bar vertical is 500 pA and horizontal is 100 ms. A2, 53/54 recorded neurons (red) with I_h co-localized with tyrosine hydroxylase immunohistochemical staining (green). A3, 1/54 recorded neurons with I_h did not co-localize with tyrosine hydroxylase. B, example neuron showing enhancement of VTA dopamine neuron firing by 10 min application of 1 μM CRF. Inset scale bar vertical is 20 mV and horizontal is 2.5 s. C, average effect of 1 μM CRF application (10 min) on firing rate of VTA dopamine neurons ($n = 14$). D, significant maximal increases in firing rate on dopamine neurons were observed with 1 μM CRF ($n = 14$) and 500 nM CRF ($n = 6$), but not with 100 nM CRF ($n = 5$). ** $P < 0.01$, *** $P < 0.001$.

control, Fig. 3A and C), or with 20 μM PKI, a direct PKA antagonist ($31.6 \pm 3.8\%$ over baseline, $n = 7$, $P > 0.05$ relative to control, Fig. 3A and C). CRF receptors also couple to the PLC–PKC signalling cascade (Blank *et al.* 2003; Ungless *et al.* 2003; Hauger *et al.* 2006), and consistent with a role for this pathway, we found that the PLC antagonist, U-73122 (1 μM), significantly attenuated the effect of CRF on dopamine cell firing ($13.4 \pm 7.8\%$ over baseline, $n = 7$, $P < 0.05$ compared to CRF alone,

Fig. 3B and C). Similarly, the general PKC antagonist, bisindomaleimide (BIS, 1 μM) also significantly prevented the CRF-dependent increase in firing ($7.7 \pm 4.2\%$ over baseline, $n = 8$, $P < 0.01$ compared to CRF alone, Fig. 3B and C). These results demonstrate that CRF activation of CRF-R1 stimulated the PLC–PKC signalling pathway to increase the firing rate in VTA dopamine neurons.

We next sought to determine the ionic target affected by CRF that mediates the increased firing rate in dopamine




Figure 2. CRF increased the firing of VTA dopamine neurons through the CRF-R1

A, the CRF-R1 agonist, oCRF (1 μM), increased the firing rate of VTA dopamine neurons ($n = 7$), while B, the CRF-R2 agonist, urocortin II (UCN II, 1 μM) did not ($n = 7$). CRF receptor antagonists were applied for 5 min prior to and during CRF exposure. C, the non-specific CRF receptor antagonist (d-Phe-CRF, 1 μM , black circles, $n = 7$) and the CRF-R1 antagonist (CP-154,526, 3 μM , red squares, $n = 7$), but not the CRF-R2 antagonist (AS-30, 250 nM, blue triangles, $n = 7$) prevented the increase in firing by CRF. D, summary of the effects of various CRF receptor agonists and antagonists on maximum changes in firing. $***P < 0.001$ from baseline firing. $##P < 0.01$, $\#P < 0.05$, respectively, reduced from CRF alone. E, CRF increased the firing of VTA dopamine neurons in CRF-R1 $^{+/+}$ mice (black circles, $n = 5$) and in CRF-R1 $^{+/-}$ mice (red squares, $n = 6$), though to a lesser degree than in CRF-R1 $^{+/+}$ mice, and did not affect firing in CRF-R1 $^{-/-}$ mice (blue triangles, $n = 4$). F, CRF enhanced the firing of VTA dopamine neurons to similar levels in CRF-R2 $^{+/+}$ (black circles, $n = 7$), CRF-R2 $^{+/-}$ (red squares, $n = 10$) animals and CRF-R2 $^{-/-}$ mice (blue triangles, $n = 5$).

neurons. The most pronounced alteration by CRF on the action potential dynamics was a significant reduction in the peak of the after-hyperpolarization potential (AHP, Fig. 1B or refer to Supplemental Fig. 3 for overlay) from -63.6 ± 0.2 mV during baseline firing to -58.6 ± 0.1 mV during CRF application ($n = 14$, $P < 0.001$). Although




Figure 3. PLC and PKC are required for CRF to increase the firing of VTA dopamine neurons

Inhibitors to intracellular signalling pathways were included in the internal recording solution. *A*, $100 \mu\text{M}$ Rp-cAMPs (red squares, $n = 8$) or $20 \mu\text{M}$ PKI (blue triangles, $n = 7$), did not prevent the effect of CRF on the firing of VTA dopamine neurons. *B*, U-73122 ($1 \mu\text{M}$, red squares, $n = 7$) and BIS ($1 \mu\text{M}$, blue triangles, $n = 8$), both blocked the increase in VTA dopamine neuron firing by CRF. *C*, summary of the effects of various intracellular signalling pathway inhibitors on the maximal change in firing by CRF. *** $P < 0.001$ relative to CRF alone. ** $P < 0.01$ relative to CRF alone.

changes in the firing rate can indirectly modulate the AHP, we first assayed whether currents that might contribute to the generation of the AHP, such as calcium-activated potassium currents ($I_{K(\text{Ca})}$) (Sah, 1996), A-type potassium currents (I_A) (Hahn *et al.* 2003), inwardly rectifying potassium currents ($I_{K\text{ir}}$) (Uchida *et al.* 2000) and I_h (Zolles *et al.* 2006), played a role in the CRF-induced enhancement of VTA dopamine neuron firing. Inhibiting calcium currents and $I_{K(\text{Ca})}$ by including 10 mM BAPTA (Tozzi *et al.* 2003) in the recording solution expectedly reduced the AHP relative to cells recorded without BAPTA in the recording solution (-63.6 ± 0.2 mV for control, $n = 14$; -56.3 ± 0.8 mV for BAPTA, $n = 8$, $P < 0.001$). However, BAPTA did not prevent CRF from increasing neuronal firing ($40.9 \pm 15.4\%$ over baseline, $n = 8$, $P > 0.05$ compared to CRF alone, Fig. 4D). Although CRF can excite CA3 pyramidal hippocampal neurons by reducing the AHP through inhibition of $I_{K(\text{Ca})}$ (Aldenhoff *et al.* 1983; Haug & Storm, 2000), this mechanism did not account for the CRF effect in VTA dopamine neurons.

I_A is present in VTA dopamine neurons and regulates the frequency of cell firing (Hahn *et al.* 2003; Koyama & Appel, 2006), making it another potential target for CRF. Blocking the slow I_A with $10 \mu\text{M}$ 4-aminopyridine (4-AP) increased neuron firing ($21.5 \pm 10.9\%$ over baseline, $n = 6$), but did not prevent a further enhancement in firing by CRF during 4-AP application ($34.8 \pm 6.3\%$ over firing rate during 4-AP application, $n = 6$, $P > 0.05$ compared to CRF alone, Fig. 4D). 4-AP at higher concentrations ($30 \mu\text{M}$, $60 \mu\text{M}$ and $100 \mu\text{M}$) similarly increased the firing and did not attenuate the effect of CRF (data not shown). With 4 mM 4-AP, a dose that eliminates both fast and slow I_A (Lien *et al.* 2002), the spontaneous firing of VTA dopamine neurons was drastically reduced and often eliminated, preventing analysis of the CRF-mediated effects on firing (data not shown). We next investigated the possibility that CRF increased the firing of dopamine neurons through inhibition of the $I_{K\text{ir}}$, as this current is modulated by CRF in corticotropes (Kuryshv *et al.* 1997). Activation of the $I_{K\text{ir}}$ reduces dopamine cell firing (Lacey *et al.* 1987; Werner *et al.* 1996; Uchida *et al.* 2000), and as expected, inhibition of $I_{K\text{ir}}$ with $100 \mu\text{M}$ barium (Ba^{2+}) increased the firing rate in VTA dopamine neurons ($114.2 \pm 27.2\%$ over baseline, $n = 8$). However, as observed under control conditions, CRF significantly increased the firing during Ba^{2+} application ($51.5 \pm 6.5\%$ over firing rate during Ba^{2+} application, $n = 8$, $P > 0.05$ compared to CRF alone, Fig. 4D). These experiments suggest that neither the slow I_A nor $I_{K\text{ir}}$ are required for CRF to increase VTA dopamine neuron firing.

I_h affects the AHP and firing rate in dopamine neurons (Neuhoff *et al.* 2002; Zolles *et al.* 2006), and can also be modulated by CRF in hypothalamic neurons (Qiu *et al.* 2005). Here, application of the I_h inhibitor ZD-7288 ($30 \mu\text{M}$) enhanced the AHP in VTA dopamine

neurons (Fig. 4A). Although $30 \mu\text{M}$ ZD-7288 decreased the firing in 6/6 neurons tested, the neurons did not always attain a stable firing rate with this high dose. Thus, in order to record from stable firing neurons and block I_h , we pre-incubated the brain slices (> 20 min) and continuously applied $1 \mu\text{M}$ ZD-7288, which was previously shown to effectively block I_h (Sato & Yamada, 2000). In these experiments, all recorded neurons lacked I_h due to the pharmacological block (data not shown), and we instead assayed for dopamine content by verifying the presence of a dopamine ($50 \mu\text{M}$)-induced hyperpolarization. When I_h was blocked, CRF did not enhance the firing rate in VTA neurons ($-0.7 \pm 3.8\%$ over baseline, $n = 5$, $P < 0.001$ relative to CRF alone, Fig. 4B–D), suggesting that CRF modulated I_h to increase the firing rate.

As our findings indicate a critical role for I_h in the excitatory action of CRF on dopamine neuron firing, we next assayed the direct effect of CRF on I_h using voltage-clamp recordings, by measuring I_h generated in

response to hyperpolarizing the neuron for 500 ms in 10 mV increments to -120 mV from our holding potential of -60 mV. CRF reversibly enhanced I_h (Fig. 5A) and the time course of this effect was remarkably similar to how CRF increased the firing rate (Fig. 1C). Additionally, CRF increased I_h over a range of voltages tested with an average enhancement of 62.8 ± 13.3 pA measured at the -120 mV voltage step ($n = 7$, Fig. 5D).

The CRF enhancement in I_h could be due to changes in the total current or a shift in the voltage dependence of activation. For experiments analysing the voltage dependence of activation for I_h we added 3 mM BAPTA to our recording solution and included 500 nM TTX, 10 mM tetraethylammonium chloride (TEA), 4 mM 4-AP and $300 \mu\text{M}$ Ba^{2+} in the aCSF. Neurons were held at -40 mV, given a hyperpolarizing voltage step for 1 s, and finally stepped to -60 mV to elicit I_h tail currents. Tail currents elicited after each hyperpolarizing step were normalized to the I_h generated from the -130 mV step. In eight neurons tested, CRF enhanced I_h (-47.2 ± 14.0 pA measured at




Figure 4. I_h is required for CRF to increase VTA dopamine neuron firing

A, overlay of action potentials from an example VTA dopamine neuron during baseline (black) and $30 \mu\text{M}$ ZD-7288 application (red) that highlights the increase of the AHP after I_h inhibition. Scale bar vertical is 30 mV and horizontal is 25 ms. Example (B) and summary of 5 neurons (C) demonstrating that I_h inhibition with continuous application of $1 \mu\text{M}$ ZD-7288 prevented the increase in firing rate by CRF in VTA dopamine neurons. D, pharmacological blockade of $I_{\text{K(Ca)}}$, I_{KIr} or slow I_{A} did not prevent the CRF-induced increase in firing. In contrast, inhibition of I_h prevented the increase in dopamine neuron firing by CRF. *** $P < 0.001$ relative to peak increase in firing by CRF alone.

the -120 mV voltage step), but there were no changes in the voltage dependence of activation, as CRF did not alter the Boltzmann sigmoidal fit values of the normalized tail current for the $V_{1/2}$ (baseline: -89.1 ± 1.5 mV; CRF: -88.4 ± 1.4 mV, Fig. 5C and D) or the slope factor (baseline: -7.5 ± 1.4 ; CRF: -7.2 ± 1.3 , Fig. 5C and D).

The CRF enhancement of dopamine neuron firing required PKC (Fig. 3B and C), and in agreement, PKC inhibition with BIS ($1 \mu\text{M}$) significantly attenuated the effect of CRF on I_h (-19.9 ± 5.6 pA measured at the -120 mV step, $n=9$, $P < 0.01$ relative to control internal, Fig. 6A and B), without altering the voltage dependence of activation (Supplemental Fig. 4A and B). The average baseline I_h measured at the -120 mV voltage

step was also significantly reduced by PKC antagonism, suggesting that PKC exerts a tonic regulation on I_h in VTA dopamine neurons (BIS: -147.1 ± 40.5 pA, $n=9$; control: -264.6 ± 33.8 pA, $n=7$, $P < 0.01$). In addition, we recapitulated the effect of CRF on I_h in VTA dopamine neurons using the PKC activator, phorbol 12,13-dibutyrate (PDBU, 500 nM), which enhanced I_h at a range of voltages tested, with a maximal increase at the -120 mV step of -35.4 ± 6.6 pA ($n=8$, Fig. 6C and D), without changing the voltage dependence of activation ($V_{1/2}$ baseline: -89.3 ± 2.0 mV; PDBU: -90.1 ± 2.3 mV, slope factor baseline: -7.9 ± 1.9 ; PDBU: -9.5 ± 2.3 , $n=7$, Fig. 6E and F). Similar to CRF, the PDBU-dependent enhancement of I_h was significantly attenuated by




Figure 5. CRF enhanced I_h without affecting the voltage dependence of activation for I_h in VTA dopamine neurons

A, hyperpolarizing voltage steps (500 ms) from a holding potential of -60 mV to -80 , -100 and -120 mV activated the slowly developing I_h (black), which was increased by CRF (red). Presented are raw traces (Aa), subtraction of traces (Ab), and the time course of this reversible effect for the step to -120 mV (Ac). Scale bar horizontal is 100 ms and vertical is 400 pA (Aa) or 100 pA (Ab). B, summary of the CRF enhancement of I_h at all voltage steps tested ($n=7$). C and D, CRF did not alter the voltage dependence of activation for I_h . C, example neuron demonstrating that CRF did not alter the tail current elicited by the offset of hyperpolarizing voltage steps (1 s) to -60 mV from -130 , -90 , -80 and -60 mV. Scale bar horizontal is 250 ms and vertical is 200 pA. D, summary of the effect of CRF on I_h tail currents in 8 neurons.

inhibiting PKC activity with 1 μM BIS in the internal solution (-16.6 ± 5.2 pA measured at the -120 mV step, $n = 8$, $P < 0.05$ compared to control internal).

Previous work demonstrated that cAMP positively modulates I_h by shifting the voltage dependence of activation (Zolles *et al.* 2006). In agreement, we observe a significant hyperpolarizing shift in the voltage dependence of activation when including 100 μM Rp-cAMPS, which inhibits cAMP-dependent processes, in the internal solution (control: -89.1 ± 1.5 mV, $n = 8$; Rp-cAMPS:

-93.4 ± 1.1 mV, $n = 8$, $F_{1,120} = 5.1$, $P < 0.05$, Fig. 7A and B), with no change in the slope factor (control: -7.5 ± 1.4 ; Rp-cAMPS: -8.9 ± 1.2). In these neurons application of CRF did not induce any further changes in the $V_{1/2}$ (-93.4 ± 1.3 mV) or the slope factor (-9.4 ± 1.4) values for the normalized tail current. However, in agreement with our firing results (Fig. 3A), blockade of cAMP signalling did not prevent the CRF-induced enhancement in I_h magnitude (-44.2 ± 5.4 pA measured at the step to -120 mV, $n = 6$, $P > 0.05$ relative to control internal,




Figure 6. CRF enhanced I_h through a PKC-dependent mechanism in VTA dopamine neurons

A, example neuron demonstrating that 1 μM BIS in the internal recording solution prevented an increase in I_h by CRF. Shown are hyperpolarizing voltage steps (500 ms) to -80 , -100 and -120 mV from a holding potential of -60 mV (Aa) and the subtraction of traces between treatments for each voltage step (Ab). Under these conditions, CRF did not affect the voltage dependence of activation for I_h (refer to Supplemental Fig. 4A and B). Scale bar horizontal is 100 ms and vertical is 400 pA (Aa) or 100 pA (Ab). B, BIS reduced the maximal effect of CRF on I_h for the voltage step measured at -120 mV ($n = 7$). $**P < 0.01$ relative to control internal. C, example neuron demonstrating that PDBU enhanced I_h . Shown are hyperpolarizing voltage steps (500 ms) to -80 , -100 and -120 mV from a holding potential of -60 mV (Ca) and the subtraction of traces between treatments for each voltage step (Cb). Scale bar horizontal is 100 ms and vertical is 400 pA (Ca) or 100 pA (Cb). D, PDBU enhanced I_h at a range of voltages tested ($n = 8$). E and F, example neuron (E) and summary of 8 neurons (F) showing that 500 nM PDBU did not change the tail current elicited by the offset of hyperpolarizing voltage steps (1 s) from -130 , -90 , -80 and -60 mV to the holding potential of -60 mV. Scale bar horizontal is 250 ms and vertical is 200 pA.

Fig. 7C and D). Taken together, our results demonstrated that CRF increased VTA dopamine neuron firing by acting on the CRF-R1 to stimulate the PLC–PKC signalling pathway, which in turn enhanced I_h without changing the voltage dependence of activation.

Discussion

Stress not only increases dopamine release in brain regions receiving dense VTA input (Abercrombie *et al.* 1989; Tidey & Miczek, 1996; Inglis & Moghaddam, 1999; McFarland *et al.* 2004), but also stimulates the release of CRF into the VTA (Wang *et al.* 2005), suggesting that CRF may directly excite the midbrain dopamine system. However, the cellular effects of CRF on VTA dopamine neurons are not well understood. In this regard, we examined the mechanism by which CRF affects VTA dopamine

neuron firing. Specifically, we found that CRF increased dopamine neuron firing by acting on the CRF-R1 to induce a PKC-dependent enhancement of I_h .

In order to study the effects of a stress-released neuropeptide on dopamine neurons, one must have a reliable method to identify dopamine neurons. The presence of I_h has been used to identify dopamine neurons in brain slice electrophysiological preparations (Grace & Onn, 1989; Cameron *et al.* 1997; Ford *et al.* 2006), but a recent study has questioned the validity of this link in the rat (Margolis *et al.* 2006). Here, we chose to use mice because an effect of CRF on NMDA receptor currents in VTA neurons had been previously described in this species (Ungless *et al.* 2003), and because of the availability of CRF receptor-deficient mice. Therefore, we re-examined whether the presence of I_h is a reliable predictor of dopamine content in




Figure 7. Inhibition of cAMP-dependent processes shifts the voltage dependence of activation for I_h , but does not prevent CRF from enhancing I_h in VTA dopamine neurons

A, examples of tail currents elicited by the offset of hyperpolarizing voltage steps (1 s) from -130, -90, -80 or -60 mV to the holding potential of -60 mV with Rp-cAMPS internal and before (baseline) or after addition of CRF. Scale bar horizontal is 250 ms and vertical is 200 pA. B, Rp-cAMPS did induce a significant hyperpolarizing shift in the baseline $V_{1/2}$ relative to recordings with the control recording. However, identical to control conditions, CRF did not further alter the voltage dependence of activation of I_h during recordings with the Rp-cAMPS internal solution. C and D, in addition, Rp-cAMPS in the internal recording solution did not prevent the CRF-mediated enhancement of I_h . C, example neuron demonstrating that Rp-cAMPS did not prevent an increase in I_h by CRF. Shown are hyperpolarizing voltage steps (500 ms) to -80, -100 and -120 mV from a holding potential of -60 mV (Ca) and the subtraction of traces between treatments for each voltage step (Cb). Scale bar horizontal is 100 ms and vertical is 400 pA (Ca) or 100 pA (Cb). D, Rp-cAMPS did not alter the maximal effect of CRF on I_h , measured at the voltage step to -120 mV relative to recordings with the control internal.

mice. In contrast to the findings from a recent study in rats (Margolis *et al.* 2006), we found that 98% of VTA neurons with I_h also contained TH, and that dopamine D2 receptor-mediated hyperpolarization accurately predicted dopamine content in mice. The discrepancy in our findings from the Margolis study could be attributed to the species utilized, or from differences in the location of recorded neurons. We recorded from VTA neurons in close proximity to the medial terminal nucleus of the accessory optic tract, where the Margolis study included neurons near the midline which can be difficult to visualize among fibre tracts using standard differential interference contrast imaging. Thus, our finding that I_h is a reliable marker of dopamine content in the mouse VTA is in agreement with prior results in rat from our laboratory (Borgland *et al.* 2006) and other studies (Grace & Onn, 1989; Ford *et al.* 2006), suggesting a higher correlation between the presence of I_h and TH immunoreactivity.

Since we could confidently identify dopamine neurons, we next observed that CRF, at doses previously shown to produce electrophysiological effects (Ungless *et al.* 2003; Kash & Winder, 2006), increased VTA dopamine neuron firing in a concentration-dependent manner. Furthermore, the results from experiments using CRF receptor agonists, CRF receptor antagonists and CRF receptor-deficient mice are parsimonious with the conclusion that CRF increased VTA dopamine neuron firing through activation of the CRF-R1. CRF receptors predominately couple to the cAMP–PKA pathway (Hauger *et al.* 2006), and it is through this pathway that CRF increases neuron firing in the locus coeruleus (Jedema & Grace, 2004) and hippocampus (Aldenhoff *et al.* 1983; Haug & Storm, 2000). Surprisingly, we found that CRF required the PLC–PKC pathway, but not the cAMP–PKA pathway, to increase VTA dopamine neuron firing.

To determine the ionic conductance affected by CRF, we hypothesized that CRF might affect a current involved with AHP generation because the firing rate increase was associated with a reduction in the AHP, and currents contributing to the firing frequency and AHP can be modulated by CRF (Aldenhoff *et al.* 1983; Kuryshv *et al.* 1997; Haug & Storm, 2000; Hahn *et al.* 2003; Qiu *et al.* 2005). However, we found that blockade of $I_{K(Ca)}$, I_{Kir} or slow I_A did not affect the ability of CRF to increase dopamine neuron firing. Instead, inhibition of I_h completely blocked the effect of CRF, suggesting a critical role for this current.

Four subunits of the hyperpolarization-activated, cyclic nucleotide-regulated cation channel (HCN) have been identified, which can mediate I_h (Frere *et al.* 2004). HCN2, HCN3 and HCN4, but not the HCN1, are present in midbrain dopamine neurons, though the exact subunit composition of functional channels is unknown in these neurons (Franz *et al.* 2000). A number of intracellular

signalling molecules can affect I_h including cAMP (Raes *et al.* 1997), phosphoinositides (Zolles *et al.* 2006; Fogle *et al.* 2007) and kinases (Zong *et al.* 2005). As we found that CRF increased VTA dopamine neuron firing in a PKC- and I_h -dependent manner, and that activation of I_h is depolarizing, we expected that CRF would enhance I_h through a similar PKC-dependent mechanism. Our voltage-clamp experiments confirmed this hypothesis and demonstrated that CRF, as well as the PKC activator, PDBU, enhanced the total I_h without affecting the voltage dependence of activation. Both the effect of CRF and PDBU on I_h was blocked by PKC antagonism, highlighting a critical role for positive modulation of I_h by PKC in VTA dopamine neurons. In addition, PKC antagonism reduced the baseline I_h , suggesting that HCN channels are under tonic positive regulation by PKC in these neurons.

Both the CRF- and PDBU-mediated increase in I_h were inhibited by PKC antagonism, clearly demonstrating that a PKC-dependent enhancement of I_h can occur in VTA dopamine neurons. However, others have reported different effects of phorbol esters on I_h in dopamine neurons. In particular, one study found that phorbol ester application reduced I_h , although these experiments were performed in the substantia nigra rather than the VTA, and the effect was found in only a small subset of neurons (Cathala & Paupardin-Tritsch, 1997). Another study in the VTA reported that higher doses of phorbol esters reduced I_h in dopamine neurons (Liu *et al.* 2003). However, the discrepancy with our results could be explained by the inhibition of the $Na^+–K^+$ pump by phorbol esters at higher doses, which could then affect a host of ionic conductances (Fisone *et al.* 1995).

It is unknown if PKC phosphorylates the HCN channels directly as with other kinases (Zong *et al.* 2005), or if PKC phosphorylates accessory proteins, which in turn have direct effects on HCN channels (Frere *et al.* 2004). Further experiments will be required to determine the exact mechanism of the PKC-dependent increase in I_h and how widespread this regulation of I_h is throughout the brain. Consistent with others, we found that the voltage dependence of activation for I_h is regulated by cAMP (Raes *et al.* 1997; Zolles *et al.* 2006); however, Rp-cAMPs did not attenuate the ability of CRF to increase the magnitude of I_h . Although additional ion currents may be affected by CRF, these potential interactions are probably not critically involved with CRF increasing the firing rate. Together, our results suggest that CRF stimulated the CRF-R1 to activate a novel PKC-dependent enhancement of I_h , which increased the firing in VTA dopamine neurons.

Increased dopamine neuron firing by a stress-released peptide, such as CRF, is consistent with numerous *in vivo* microdialysis studies demonstrating elevated dopamine release by stress (Abercrombie *et al.* 1989; Tidey & Miczek, 1996; Inglis & Moghaddam, 1999; McFarland *et al.* 2004). Our findings provide a cellular mechanism by which CRF

could affect dopamine-related behaviours. For example, CRF administered to the ventricles (Contarino *et al.* 2000) or directly into the VTA (Kalivas *et al.* 1987) increases locomotor activity, and the effect of ventricular infusions of CRF on motor activity is absent in CRF-R1^{-/-} mice (Contarino *et al.* 2000). In addition, our findings provide a possible mechanism by which rodents motivate to escape from stressful and threatening situations, as enhancing dopamine levels (Blanchard *et al.* 1999) and ventricular infusions of the CRF-R1 agonists (Yang *et al.* 2006) increase the speed and number of escapes from a predator, while CRF-R1 antagonists have the opposite effect (Griebel *et al.* 1998). Finally, both stress and dopamine can enhance performance on specific behaviours depending on the context (Rose, 1987; Blanchard *et al.* 1999; Cagniard *et al.* 2006), which could involve the actions of CRF in the mesolimbic system (Pecina *et al.* 2006). Therefore, these described behaviours may arise in part due to the cellular effects of CRF on VTA dopamine neurons that we elucidated in the current study. It should also be noted that the CRF has important effects in other brain structures involved with arousal and stress responses, including the locus coeruleus (Koob, 1999; Jedema *et al.* 2001; Jedema & Grace, 2004) and the amygdala (Sajdyk *et al.* 1999; Liu *et al.* 2005; Winsky-Sommerer *et al.* 2005), which may also play a significant role in the above-described behaviours.

In conclusion, we demonstrated that CRF stimulated the CRF-R1 on VTA dopamine neurons and activated a PKC-dependent enhancement of I_h , which led to increased cell firing. These results provide a potential physiological mechanism underlying the interaction between stress, dopamine and motivation, which is important for many behaviours including escape from threatening stimuli (Blanchard *et al.* 2003). Furthermore, this study identifies a link between dopamine and CRF, which together have been implicated in psychiatric disorders such as depression (Banki *et al.* 1987; Valdez, 2006; Gershon *et al.* 2007), drug abuse (McFarland *et al.* 2004; Wang *et al.* 2005; Funk *et al.* 2007) and schizophrenia (Banki *et al.* 1987; Beninger, 2006), and in this regard, highlight potential therapeutic targets to treat these disorders.

References

- Abercrombie ED, Keefe KA, DiFrischia DS & Zigmond MJ (1989). Differential effect of stress on in vivo dopamine release in striatum, nucleus accumbens, and medial frontal cortex. *J Neurochem* **52**, 1655–1658.
- Aldenhoff JB, Gruol DL, Rivier J, Vale W & Siggins GR (1983). Corticotropin releasing factor decreases postburst hyperpolarizations and excites hippocampal neurons. *Science* **221**, 875–877.
- Anstrom KK & Woodward DJ (2005). Restraint increases dopaminergic burst firing in awake rats. *Neuropsychopharmacology* **30**, 1832–1840.
- Banki CM, Bissette G, Arato M, O'Connor L & Nemeroff CB (1987). CSF corticotropin-releasing factor-like immunoreactivity in depression and schizophrenia. *Am J Psychiatry* **144**, 873–877.
- Beninger RJ (1983). The role of dopamine in locomotor activity and learning. *Brain Res* **287**, 173–196.
- Beninger RJ (2006). Dopamine and incentive learning: a framework for considering antipsychotic medication effects. *Neurotox Res* **10**, 199–209.
- Blanchard DC, Griebel G & Blanchard RJ (2003). The Mouse Defense Test Battery: pharmacological and behavioral assays for anxiety and panic. *Eur J Pharmacol* **463**, 97–116.
- Blanchard RJ, Kaawaloa JN, Hebert MA & Blanchard DC (1999). Cocaine produces panic-like flight responses in mice in the mouse defense test battery. *Pharmacol Biochem Behav* **64**, 523–528.
- Blank T, Nijholt I, Grammatopoulos DK, Randevara HS, Hillhouse EW & Spiess J (2003). Corticotropin-releasing factor receptors couple to multiple G-proteins to activate diverse intracellular signaling pathways in mouse hippocampus: role in neuronal excitability and associative learning. *J Neurosci* **23**, 700–707.
- Borgland SL, Taha SA, Sarti F, Fields HL & Bonci A (2006). Orexin A in the VTA is critical for the induction of synaptic plasticity and behavioral sensitization to cocaine. *Neuron* **49**, 589–601.
- Cagniard B, Balsam PD, Brunner D & Zhuang X (2006). Mice with chronically elevated dopamine exhibit enhanced motivation, but not learning, for a food reward. *Neuropsychopharmacology* **31**, 1362–1370.
- Cameron DL, Wessendorf MW & Williams JT (1997). A subset of ventral tegmental area neurons is inhibited by dopamine, 5-hydroxytryptamine and opioids. *Neuroscience* **77**, 155–166.
- Cathala L & Paupardin-Tritsch D (1997). Neurotensin inhibition of the hyperpolarization-activated cation current (I_h) in the rat substantia nigra pars compacta implicates the protein kinase C pathway. *J Physiol* **503**, 87–97.
- Contarino A, Dellu F, Koob GF, Smith GW, Lee KF, Vale WW & Gold LH (2000). Dissociation of locomotor activation and suppression of food intake induced by CRF in CRFR1-deficient mice. *Endocrinology* **141**, 2698–2702.
- Fisone G, Snyder GL, Fryckstedt J, Caplan MJ, Aperia A & Greengard P (1995). Na⁺,K⁺-ATPase in the choroid plexus. Regulation by serotonin/protein kinase C pathway. *J Biol Chem* **270**, 2427–2430.
- Fogle KJ, Lyashchenko AK, Turbendian HK & Tibbs GR (2007). HCN pacemaker channel activation is controlled by acidic lipids downstream of diacylglycerol kinase and phospholipase A2. *J Neurosci* **27**, 2802–2814.
- Ford CP, Mark GP & Williams JT (2006). Properties and opioid inhibition of mesolimbic dopamine neurons vary according to target location. *J Neurosci* **26**, 2788–2797.
- Franz O, Liss B, Neu A & Roeper J (2000). Single-cell mRNA expression of HCN1 correlates with a fast gating phenotype of hyperpolarization-activated cyclic nucleotide-gated ion channels (I_h) in central neurons. *Eur J Neurosci* **12**, 2685–2693.
- Frere SG, Kuisle M & Luthi A (2004). Regulation of recombinant and native hyperpolarization-activated cation channels. *Mol Neurobiol* **30**, 279–305.

- Funk CK, Zorrilla EP, Lee MJ, Rice KC & Koob GF (2007). Corticotropin-releasing factor 1 antagonists selectively reduce ethanol self-administration in ethanol-dependent rats. *Biol Psychiatry* **61**, 78–86.
- Gershon AA, Vishne T & Grunhaus L (2007). Dopamine D2-like receptors and the antidepressant response. *Biol Psychiatry* **61**, 145–153.
- Grace AA & Onn SP (1989). Morphology and electrophysiological properties of immunocytochemically identified rat dopamine neurons recorded *in vitro*. *J Neurosci* **9**, 3463–3481.
- Griebel G, Perrault G & Sanger DJ (1998). Characterization of the behavioral profile of the non-peptide CRF receptor antagonist CP-154,526 in anxiety models in rodents. Comparison with diazepam and buspirone. *Psychopharmacology (Berl)* **138**, 55–66.
- Hahn J, Tse TE & Levitan ES (2003). Long-term K⁺ channel-mediated dampening of dopamine neuron excitability by the antipsychotic drug haloperidol. *J Neurosci* **23**, 10859–10866.
- Haug T & Storm JF (2000). Protein kinase A mediates the modulation of the slow Ca²⁺-dependent K⁺ current, I_{sAHP}, by the neuropeptides CRF, VIP, and CGRP in hippocampal pyramidal neurons. *J Neurophysiol* **83**, 2071–2079.
- Hauger RL, Risbrough V, Brauns O & Dautzenberg FM (2006). Corticotropin releasing factor (CRF) receptor signaling in the central nervous system: new molecular targets. *CNS Neurol Disord Drug Targets* **5**, 453–479.
- Horger BA & Roth RH (1996). The role of mesoprefrontal dopamine neurons in stress. *Crit Rev Neurobiol* **10**, 395–418.
- Inglis FM & Moghaddam B (1999). Dopaminergic innervation of the amygdala is highly responsive to stress. *J Neurochem* **72**, 1088–1094.
- Jedema HP, Finlay JM, Sved AF & Grace AA (2001). Chronic cold exposure potentiates CRH-evoked increases in electrophysiological activity of locus coeruleus neurons. *Biol Psychiatry* **49**, 351–359.
- Jedema HP & Grace AA (2004). Corticotropin-releasing hormone directly activates noradrenergic neurons of the locus ceruleus recorded *in vitro*. *J Neurosci* **24**, 9703–9713.
- Johnson SW & North RA (1992). Two types of neuron in the rat ventral tegmental area and their synaptic inputs. *J Physiol* **450**, 455–468.
- Kalivas PW, Duffy P & Latimer LG (1987). Neurochemical and behavioral effects of corticotropin-releasing factor in the ventral tegmental area of the rat. *J Pharmacol Exp Ther* **242**, 757–763.
- Kash TL & Winder DG (2006). Neuropeptide Y and corticotropin-releasing factor bi-directionally modulate inhibitory synaptic transmission in the bed nucleus of the stria terminalis. *Neuropharmacology* **51**, 1013–1022.
- Kauer JA (2004). Learning mechanisms in addiction: synaptic plasticity in the ventral tegmental area as a result of exposure to drugs of abuse. *Annu Rev Physiol* **66**, 447–475.
- Koob GF (1999). Corticotropin-releasing factor, norepinephrine, and stress. *Biol Psychiatry* **46**, 1167–1180.
- Korotkova TM, Brown RE, Sergeeva OA, Ponomarenko AA & Haas HL (2006). Effects of arousal- and feeding-related neuropeptides on dopaminergic and GABAergic neurons in the ventral tegmental area of the rat. *Eur J Neurosci* **23**, 2677–2685.
- Koyama S & Appel SB (2006). A-type K⁺ current of dopamine and GABA neurons in the ventral tegmental area. *J Neurophysiol* **96**, 544–554.
- Kuryshv YA, Haak L, Childs GV & Ritchie AK (1997). Corticotropin releasing hormone inhibits an inwardly rectifying potassium current in rat corticotropes. *J Physiol* **502**, 265–279.
- Lacey MG, Mercuri NB & North RA (1987). Dopamine acts on D2 receptors to increase potassium conductance in neurones of the rat substantia nigra zona compacta. *J Physiol* **392**, 397–416.
- Lavicky J & Dunn AJ (1993). Corticotropin-releasing factor stimulates catecholamine release in hypothalamus and prefrontal cortex in freely moving rats as assessed by microdialysis. *J Neurochem* **60**, 602–612.
- Lien CC, Martina M, Schultz JH, Ehmke H & Jonas P (2002). Gating, modulation and subunit composition of voltage-gated K⁺ channels in dendritic inhibitory interneurons of rat hippocampus. *J Physiol* **538**, 405–419.
- Liu J, Yu B, Orozco-Cabal L, Grigoriadis DE, Rivier J, Vale WW, Shinnick-Gallagher P & Gallagher JP (2005). Chronic cocaine administration switches corticotropin-releasing factor₂ receptor-mediated depression to facilitation of glutamatergic transmission in the lateral septum. *J Neurosci* **25**, 577–583.
- Liu Z, Bunney EB, Appel SB & Brodie MS (2003). Serotonin reduces the hyperpolarization-activated current (I_h) in ventral tegmental area dopamine neurons: involvement of 5-HT₂ receptors and protein kinase C. *J Neurophysiol* **90**, 3201–3212.
- Lodge DJ & Grace AA (2005). Acute and chronic corticotropin-releasing factor 1 receptor blockade inhibits cocaine-induced dopamine release: correlation with dopamine neuron activity. *J Pharmacol Exp Ther* **314**, 201–206.
- McFarland K, Davidge SB, Lapish CC & Kalivas PW (2004). Limbic and motor circuitry underlying footshock-induced reinstatement of cocaine-seeking behavior. *J Neurosci* **24**, 1551–1560.
- Magee JC (1998). Dendritic hyperpolarization-activated currents modify the integrative properties of hippocampal CA1 pyramidal neurons. *J Neurosci* **18**, 7613–7624.
- Margolis EB, Lock H, Hjelmstad GO & Fields HL (2006). The ventral tegmental area revisited: is there an electrophysiological marker for dopaminergic neurons? *J Physiol* **577**, 907–924.
- Neuhoff H, Neu A, Liss B & Roeper J (2002). I_h channels contribute to the different functional properties of identified dopaminergic subpopulations in the midbrain. *J Neurosci* **22**, 1290–1302.
- Pecina S, Schulkin J & Berridge KC (2006). Nucleus accumbens corticotropin-releasing factor increases cue-triggered motivation for sucrose reward: paradoxical positive incentive effects in stress? *BMC Biol* **4**, 8.

- Phillips PE, Walton ME & Jhou TC (2007). Calculating utility: preclinical evidence for cost-benefit analysis by mesolimbic dopamine. *Psychopharmacology (Berl)* **191**, 483–495.
- Qiu DL, Chu CP, Shirasaka T, Tsukino H, Nakao H, Kato K, Kunitake T, Katoh T & Kannan H (2005). Corticotrophin-releasing factor augments the I_H in rat hypothalamic paraventricular nucleus parvocellular neurons *in vitro*. *J Neurophysiol* **94**, 226–234.
- Raes A, Wang Z, van den Berg RJ, Goethals M, Van de Vijver G & van Bogaert PP (1997). Effect of cAMP and ATP on the hyperpolarization-activated current in mouse dorsal root ganglion neurons. *Pflugers Arch* **434**, 543–550.
- Rose RM (1987). Neuroendocrine effects of work stress. In *Work Stress: Health Care Systems in the Workplace*, ed. Quick JC, Bhagal RS, Dalton JE & Quick JD, pp. 130–147. Praeger, New York.
- Sah P (1996). Ca^{2+} -activated K^+ currents in neurones: types, physiological roles and modulation. *Trends Neurosci* **19**, 150–154.
- Sajdyk TJ, Schober DA, Gehlert DR & Shekhar A (1999). Role of corticotropin-releasing factor and urocortin within the basolateral amygdala of rats in anxiety and panic responses. *Behav Brain Res* **100**, 207–215.
- Salamone JD (1996). The behavioral neurochemistry of motivation: methodological and conceptual issues in studies of the dynamic activity of nucleus accumbens dopamine. *J Neurosci Meth* **64**, 137–149.
- Sarnyai Z, Shaham Y & Heinrichs SC (2001). The role of corticotropin-releasing factor in drug addiction. *Pharmacol Rev* **53**, 209–243.
- Satoh TO & Yamada M (2000). A bradycardiac agent ZD7288 blocks the hyperpolarization-activated current (I_h) in retinal rod photoreceptors. *Neuropharmacology* **39**, 1284–1291.
- Tan H, Zhong P & Yan Z (2004). Corticotropin-releasing factor and acute stress prolongs serotonergic regulation of GABA transmission in prefrontal cortical pyramidal neurons. *J Neurosci* **24**, 5000–5008.
- Tidey JW & Miczek KA (1996). Social defeat stress selectively alters mesocorticolimbic dopamine release: an *in vivo* microdialysis study. *Brain Res* **721**, 140–149.
- Tozzi A, Bengtson CP, Longone P, Carignani C, Fusco FR, Bernardi G & Mercuri NB (2003). Involvement of transient receptor potential-like channels in responses to mGluR-I activation in midbrain dopamine neurons. *Eur J Neurosci* **18**, 2133–2145.
- Uchida S, Akaike N & Nabekura J (2000). Dopamine activates inward rectifier K^+ channel in acutely dissociated rat substantia nigra neurones. *Neuropharmacology* **39**, 191–201.
- Ungless MA, Singh V, Crowder TL, Yaka R, Ron D & Bonci A (2003). Corticotropin-releasing factor requires CRF binding protein to potentiate NMDA receptors via CRF receptor 2 in dopamine neurons. *Neuron* **39**, 401–407.
- Valdez GR (2006). Development of CRF1 receptor antagonists as antidepressants and anxiolytics: progress to date. *CNS Drugs* **20**, 887–896.
- Van Pett K, Viau V, Bittencourt JC, Chan RK, Li HY, Arias C, Prins GS, Perrin M, Vale W & Sawchenko PE (2000). Distribution of mRNAs encoding CRF receptors in brain and pituitary of rat and mouse. *J Comp Neurol* **428**, 191–212.
- Wang B, Shaham Y, Zitzman D, Azari S, Wise RA & You ZB (2005). Cocaine experience establishes control of midbrain glutamate and dopamine by corticotropin-releasing factor: a role in stress-induced relapse to drug seeking. *J Neurosci* **25**, 5389–5396.
- Werner P, Hussy N, Buell G, Jones KA & North RA (1996). D2, D3, and D4 dopamine receptors couple to G protein-regulated potassium channels in *Xenopus* oocytes. *Mol Pharmacol* **49**, 656–661.
- Winsky-Sommerer R, Boutrel B & de Lecea L (2005). Stress and arousal: the corticotropin-releasing factor/hypocretin circuitry. *Mol Neurobiol* **32**, 285–294.
- Wise RA (2004). Dopamine, learning and motivation. *Nat Rev Neurosci* **5**, 483–494.
- Yang M, Farrokhi C, Vasconcellos A, Blanchard RJ & Blanchard DC (2006). Central infusion of ovine CRF (oCRF) potentiates defensive behaviors in CD-1 mice in the Mouse Defense Test Battery (MDTB). *Behav Brain Res* **171**, 1–8.
- Zhou QY & Palmiter RD (1995). Dopamine-deficient mice are severely hypoactive, adipsic, and aphagic. *Cell* **83**, 1197–1209.
- Zolles G, Klocker N, Wenzel D, Weisser-Thomas J, Fleischmann BK, Roeper J & Fakler B (2006). Pacemaking by HCN channels requires interaction with phosphoinositides. *Neuron* **52**, 1027–1036.
- Zong X, Eckert C, Yuan H, Wahl-Schott C, Abicht H, Fang L, Li R, Mistrik P, Gerstner A, Much B, Baumann L, Michalakakis S, Zeng R, Chen Z & Biel M (2005). A novel mechanism of modulation of hyperpolarization-activated cyclic nucleotide-gated channels by Src kinase. *J Biol Chem* **280**, 34224–34232.

Acknowledgements

This work was supported by the State of California for medical research on alcohol and substance abuse through the University of California, San Francisco (A.B.) and the National Institute on Drug Abuse R01 DA016782-04 (A.B.), F31 DA21464-01 (M.J.W.). We thank Wylie Vale for the CRF receptor-deficient mice, Ling Wang for her assistance in histology, Viktor Kharazia for his assistance in immunocytochemistry, Lisa G. Daitch for proofreading the manuscript, as well as the members of the Bonci and Phillips labs for their critical input.

Supplemental material

Online supplemental material for this paper can be accessed at: <http://jp.physoc.org/cgi/content/full/jphysiol.2007.150078/DC1> and <http://www.blackwell-synergy.com/doi/suppl/10.1113/jphysiol.2007.150078>